

WIENER PHYSIKALISCHES KOLLOQUIUM

TU-WIEN - UNIVERSITÄT WIEN

SS 2015

Einladung zum Vortrag von

Phil Bucksbaum

Department of Physics
Stanford University USA

Investigating ultrafast dynamics in atoms and molecules with strong short wavelength coherent radiation

The time scale for internal motion in atoms and small molecules is determined by their Angstrom sizes and Rydberg binding energies to be femtoseconds or shorter. The binding fields for the outermost electrons are tens of volts per Angstrom, while inner electrons can be bound by kilovolts or more. I will describe recent experiments designed to study the interaction of atoms and molecules utilizing laser fields on these scales of time, photon energy and field strength. Two kinds of laser sources are employed: Strong focused infrared lasers create these extreme conditions within a single optical cycle, and thereby induce atomic phenomena that evolve during fractions of a femtosecond. This is the regime of high harmonic generation and above-threshold ionization. X-ray free electron lasers can also produce these extreme fields, but at much higher oscillation frequencies. This is the regime of rapid inner shell ionization and Auger relaxation. Both types of strong-field phenomena induce dynamics on femtosecond or faster time scales. Our experiments seek to track this motion and reveal the underlying internal mechanisms responsible for them.

22. Juni 2015, 17:30 hrs

TU Wien-Freihaus, Hörsaal 5, 2. Stock, grüner Bereich